

PATIENT DEMOGRAPHICS

Patient Name: _____ DOB: _____ Phone: _____
 Address: _____ City/ST/Zip: _____
 Allergies: _____ NKDA Weight: _____ lbs kg Height: _____ in cm
Patient Status: New to Therapy Dose or Frequency Change Order Renewal

INSURANCE INFORMATION: Please attach copy of insurance card (front and back).

DIAGNOSIS*

***ICD 10 Code Required**

- Idiopathic chronic gout (M1A.00-M1A.09), ICD10 _____
- Lead-induced chronic gout (M1A.10-M1A.19), ICD10 _____
- Drug-induced chronic gout (M1A.20-M1A.29), ICD10 _____
- Other: _____, ICD10 _____
- Chronic gout due to renal impairment (M1A.30-M1A.39), ICD10 _____
- Other secondary chronic gout (M1A.40-M1A.49), ICD10 _____
- Chronic gout, unspecified (M1A.9XX0-M1A.9XX1), ICD10 _____

INFUSION ORDERS

MEDICATION	DOSE	DIRECTIONS/DURATION
Krystexxa® (Pegloticase)	8 mg	Infuse IV over 2 hours once every 2 weeks x 1 year <input type="checkbox"/> Notify physician if uric acid >6 mg/mL before infusing <input type="checkbox"/> Observe patient for 1 hour after completion of infusion

Is patient currently receiving therapy above from another facility?

NO YES

If yes, Facility Name: _____

Date of last treatment: _____ Date of next treatment: _____

PRE-MEDICATION ORDERS

- Acetaminophen 650mg PO 30 minutes prior to infusion
- Diphenhydramine 25mg/50mg PO 30 minutes prior to infusion
- Methylprednisolone 100mg IV 30 minutes prior to infusion
- Other: _____

LAB ORDERS

- Labs to be drawn by:** Infusion Center Referring Physician
- Serum uric acid – baseline and prior to each infusion with results
 - Other: _____

REFERRING PHYSICIAN INFORMATION

Physician Signature: _____ Date: _____
 Physician Name: _____ NPI: _____ TIN: _____ Specialty: _____
 Address: _____ City/ST/Zip: _____
 Contact Person: _____ Phone #: _____ Fax #: _____
 Email Where Follow Up Documentation Should Be Sent: _____

REQUIRED CLINICAL DOCUMENTATION

Please attach medical records: Initial H&P, current MD progress notes, medication list, and labs/test results to support diagnosis.

Clinical Information, select all that apply:

- The patient has refractory chronic gout as evidenced by the following:
 - History of at least 2 gout flares in the previous 12 months
 - At least 1 gout tophus
 - Gouty arthritis
 - Other: _____
- The patient has a baseline serum uric acid of ≥ 6 mg/dL prior to initiating Krystexxa.
- The patient does not have Glucose-6-phosphate dehydrogenase (G6PD) deficiency.
- Krystexxa will be co-administered with weekly oral methotrexate unless there is a documented contraindication.

LAB and TEST RESULTS (required)

- Provide copy of baseline serum uric acid level
- Provide copy of Glucose-6-phosphate dehydrogenase (G6PD) test

PRIOR FAILED THERAPIES FOR CHRONIC GOUT

Medication Failed: _____ Dates of Treatment: _____ Reason for D/C: _____
 Medication Failed: _____ Dates of Treatment: _____ Reason for D/C: _____
 Medication Failed: _____ Dates of Treatment: _____ Reason for D/C: _____
 Medication Failed: _____ Dates of Treatment: _____ Reason for D/C: _____
 Medication Failed: _____ Dates of Treatment: _____ Reason for D/C: _____